

Available online at www.sciencedirect.com



Tectonophysics xx (2004) xxx-xxx

TECTONOPHYSICS

www.elsevier.com/locate/tecto

Production and recycling of oceanic crust in the early Earth

P. van Thienen*, A.P. van den Berg, N.J. Vlaar

Department of Theoretical Geophysics, Institute of Earth Sciences, University of Utrecht, P.O. Box 80.021, 3508 TA, Utrecht, The Netherlands

Received 18 June 2003; accepted 27 April 2004

Abstract

Because of the strongly different conditions in the mantle of the early Earth regarding temperature and viscosity, present-day geodynamics cannot simply be extrapolated back to the early history of the Earth. We use numerical thermochemical convection models including partial melting and a simple mechanism for melt segregation and oceanic crust production to investigate an alternative suite of dynamics which may have been in operation in the early Earth. Our modelling results show three processes that may have played an important role in the production and recycling of oceanic crust: (1) Small-scale ($x \times 100$ km) convection involving the lower crust and shallow upper mantle. Partial melting and thus crustal production takes place in the upwelling limb and delamination of the eclogitic lower crust in the downwelling limb. (2) Large-scale resurfacing events in which (nearly) the complete crust sinks into the (eventually lower) mantle, thereby forming a stable reservoir enriched in incompatible elements in the deep mantle. New crust is simultaneously formed at the surface from segregating melt. (3) Intrusion of lower mantle diapirs with a high excess temperature (about 250 K) into the upper mantle, causing massive melting and crustal growth. This allows for plumes in the Archean upper mantle with a much higher excess temperature than previously expected from theoretical considerations.

© 2004 Elsevier B.V. All rights reserved.

Keywords: Mantle convection; Numerical modelling; Archean; Oceanic crustal growth; Crustal recycling; Mantle chemistry

1. Introduction

As no oceanic crust on Earth is older than about 200 million years, we have no direct evidence on the nature of oceanic crust in the early Earth. Obducted

oceanic sequences in the form of ophiolites, in a strict definition, go back to about 800–900 million years before present, or back to about 2000 My when a broader definition is applied (Helmstaedt and Scott, 1992). The Archean, however, lacks ophiolites (Hamilton, 1998). A recent find of a 2.505-Gy ophiolite sequence in North China (Kusky et al., 2001) has been disputed by others (Zhai et al., 2002).

From theoretical considerations, several characteristics can be inferred about oceanic lithosphere in the early Earth. As a consequence of the higher mantle

^{*} Corresponding author. Present address: Institut de Physique du Globe de Paris, 4 Avenue de Neptune, Saint-Maur-des-Fossés, France.

E-mail addresses: thienen@ipgp.jussieu.fr (P. van Thienen), berg@geo.uu.nl (A.P. van den Berg), vlaar@geo.uu.nl (N.J. Vlaar).

^{0040-1951/}\$ - see front matter © 2004 Elsevier B.V. All rights reserved. doi:10.1016/j.tecto.2004.04.027

temperature, partial melting in shallow convective upwellings produced a layering of basaltic crust and underlying depleted (lherzolitic-harzburgitic) mantle peridotite much thicker than expected under modern day mid-oceanic ridges. As the extent of the continents is thought to have progressively grown through the Earth's history (McCulloch and Bennett, 1994), the extent of the complementary part which is the oceanic domain must have been larger than it is today. The more extensive differentiation of mantle peridotite generates more positive chemical buoyancy which takes a much longer time to be balanced by negative buoyancy due to cooling, and which would therefore be much more difficult to subduct on reasonable time scales (Sleep and Windley, 1982; Vlaar, 1985, 1986; Vlaar and Van den Berg, 1991). The dynamics of an oceanic lithosphere in the early Earth must therefore have been different from those in the present Earth. Crustal recycling must however have taken place because of the present relative paucity of Archean crustal material. Furthermore, isotopic evidence indicates crustal recycling during the Archean (Nielsen et al., 2002).

An alternative mechanism of recycling of oceanic crust into the mantle was proposed by Vlaar et al. (1994). When a basaltic crustal layer becomes sufficiently thick, a phase transition to eclogite may occur in the deeper parts (e.g. Hacker, 1996). In the presence of a sufficiently weak rheology, delamination of this dense crustal layer may take place and thus recycling of dense eclogite into the upper mantle. The space vacated by the delaminating eclogite can be filled up by inflowing mantle material, which may be fertile and undergo decompression melting, thus adding new material to the crust. This recycling mechanism may have contributed significantly to the early cooling of the Earth during the Archean (Vlaar et al., 1994). However, as the fertile mantle material replacing delaminated eclogite only undergoes partial melting, it produces a smaller volume of crust compared to the crust involved in the delamination. Therefore, the mechanism cannot be self-sustaining, since there is no complete recovery of the crust. An additional source of crustal material in the form of an active upwelling of fertile mantle rock (for example in mantle diapirs) is required to keep this process going. Related models concerning the delamination of eclogite have been presented by Parmentier and Hess

(1992) in a general and abstract sense for terrestrial planets and by Zegers and Van Keken (2001) for the more specific subject of generation of continental material.

Geochemistry may give insight into the chemical state of the early Earth. Differentiation may be expected both in a solidifying magma ocean and during partial melting of an already solidified mantle.

The chemistry of the earliest material found on the present Earth's surface, 4.0–4.4-Gy detrital zircons (Wilde et al., 2001; Peck et al., 2001), indicates growth in a granitic magma, which means that production of continental crustal material and thus differentiation was already taking place.

The early mantle was already (partially) depleted in incompatible elements (as shown by Nd and Hf isotopic evidence, Hamilton et al., 1983; Patchett, 1983; Vervoort et al., 1996; Blichert-Toft et al., 1999), with a minimum age estimate of the derivation of the depleted mantle from a uniform chondritic reservoir of 4.08 Gy (Amelin et al., 2000). An increasing trend with decreasing age in the degree of depletion of the mantle from which the crustal rocks derived has been observed (Amelin et al., 2000).

Differentiation of the proto-mantle in a partially molten regime may be expected, but the result is only a change in the relative proportions of compatible and incompatible elements, retaining more or less chondritic relative abundances for the major elements (Abe, 1993). Computations by Abe (1997) show that differentiation in a terrestrial magma ocean spanning at least the upper mantle probably took place and may be related to the Nd and Hf isotopic evidence for early mantle depletion.

Calculations of fractionation upon solidification of a terrestrial magma ocean of lithophile element ratios indicate no surviving evidence of mineral fractionation (McFarlane et al., 1994), though element partitioning data may not fully constrain this (Presnall et al., 1998).

These combined results from geochemistry suggest that chemical differentiation in the early mantle was an important process, as seen from isotope ratios, but that large-scale inhomogeneities may have been rapidly remixed, as suggested by lithophile element ratios. This makes it difficult to come up with a viable starting scenario for numerical models of the mantle in the early Earth.

Numerical mantle convection models starting from a pristine mantle show the development of chemical heterogeneity in the mantle. Depending on the rate of internal heating due to the decay of radioactive isotopes, stratification of depleted and undepleted mantle material or mixing of these may occur on whole mantle scale (Ogawa, 1988, 2000; Kameyama et al., 1996). Walzer and Hendel (1997) modelled a simplified chemical segregation in the mantle, generating continents. In a more complex setup, they modelled the development of distinct pristine lower mantle and depleted upper and uppermost lower mantle reservoirs (Walzer and Hendel, 1999). The development of chemical stratification in the upper mantle underneath continents was demonstrated in thermochemical convection models (De Smet et al., 1998, 2000; De Smet, 1999). In these models, a continental root of depleted peridotite of about 200 km develops, underneath which the remaining part of the upper mantle remains relatively fertile. Similar models by Schott et al. (2001) applied to Mars produce a thick (500 km) buoyant layer of melt residue, inhibiting recycling of crustal material.

Here, we present results of numerical modelling experiments of mantle convection including pressure release partial melting. The model includes a simple approximate melt segregation mechanism and basalt to eclogite phase transition, to account for the dynamic accumulation and recycling of the crust in an upper mantle subject to secular cooling. The focus of our investigations is on the dynamics of the production of oceanic crust and its recycling through the eclogite phase into the mantle. Production and recycling of basaltic crust through the eclogite phase has been studied before by Dupeyrat and Sotin (1995) in isoviscous models applied to Venus. Our approach is innovative in the sense that we use a 2-D convection model which combines mantle melting and crustal production through approximate melt segregation with phase transitions in both the mantle and crustal phases and a composite rheology model including yielding. Specifically, the composite rheology model has a strong impact on the dynamics compared to an isoviscous case. The current model allows us to study crustal production and recycling as it is generated self-consistently by the model. We investigate the conditions in favor of such a crustal recycling mechanism and discuss its consequences for the cooling history of the Earth. We also investigate the development and sustainability of the chemical differentiation which is caused by the process, both concerning the major elements and incompatible trace elements.

2. Numerical model setup

2.1. Description of the numerical model

We have used a 2-D Cartesian thermochemical convection code including partial melting. An extended Boussinesq approach was used, assuming infinite Prandtl number, including viscous dissipation, adiabatic compression, and latent heat of melting. Partial melting is modelled as an (irreversible) increase in the degree of depletion F, which has been defined here as the mass fraction of melt that is extracted from an initially unmelted material control volume of mantle material. The melt that is produced is extracted instantaneously and deposited at the top boundary. Here, the melt flux is transformed into an inflow boundary condition, thus producing a basaltic crust. In the current approximation, the compaction of the residual mantle due to this melt removal is neglected. This results in an overestimate of the volume of depleted peridotite. In most cases in our experiments, the degree of depletion is around 0.10 or less, and the effect is only minor. In some regions, the degree of depletion may be several tens of percents. But because these regions are generally small, we expect that ignoring compaction has little influence on the overall dynamics. Time-dependent internal heating due to the decay of radioactive elements was included and coupled to the concentration of a single incompatible element, which has an initial concentration of 1 and fractionates upon partial melting with a distribution coefficient of 0.01 (see Section 2.3). Undepleted mantle material is characterized by a time-dependent heat productivity with an initial value of 15×10^{-12} W/kg and a half-life value of radioactive decay of 2.5 Gy. For the compositionally evolving mantle material, this time-dependent value is multiplied with the local trace element concentration. The extraction of heat producing elements during partial melting results in a concentration of the heat productivity in the basalt produced by the partial melting.

The model is described by the following nondimensional equations (De Smet et al., 1998). The energy equation (see Table 1 for explanation of the symbols used):

$$\frac{\mathrm{d}T}{\mathrm{d}t} - Di(T+T_0)w
= \nabla^2 T - \frac{\Delta S}{c_{\mathrm{p}}} \frac{\mathrm{d}F}{\mathrm{d}t}(T+T_0)
+ \sum_k \gamma_k \frac{Rc_k}{Ra} Di(T+T_0) \frac{\mathrm{d}\Gamma_k}{\mathrm{d}t} + \frac{Di}{Ra} \Phi + R_{\mathrm{H}}H$$
(1)

The heating due to viscous dissipation is set to zero in the calculations to avoid extreme heating rates in the upper crust, where laterally varying amounts of crustal material is forced into the system at considerable rates, see below. We have verified that this has a minor effect on the resulting dynamics and thermal development of the subcrustal domain.

We use the Stokes equation in combination with a continuity equation for an infinite Prandtl number incompressible fluid:

$$\nabla \left[\eta(p, T, \tau, F) \left(\nabla \vec{u} + \nabla \vec{u}^{T} \right) \right] - \nabla \Delta p$$
$$= \left(RaT + RbF - \sum_{k} Rc_{k}\Gamma_{k} \right) \hat{z}$$
(2)

$$\nabla \cdot \vec{u} = 0 \tag{3}$$

The formulation of the variable viscosity term of the nonlinear rheological model will be described in Section 2.2. Evolution of the degree of depletion is described by the following equation:

$$\frac{\mathrm{d}F}{\mathrm{d}t} = S(P, T, F) \tag{4}$$

The source function S in Eq. (4) describes the distribution of partial melting, applying a simple parameterization of the melting phase diagram of mantle peridotite (De Smet et al., 1998). We use a third-order polynomial parameterizations of the solidus and the liquidus, based on Herzberg and Zhang (1996), down to a depth of 400 km, assuming that melt produced below this depth is not segregated. Our isobaric melting curve, which is based on data presented by Jaques and Green (1980), is linear (see De Smet et al., 1998).

The set of Eqs. (1)-(4) was solved using finite element methods and a Predictor-Corrector time integration scheme, described in Van den Berg et al. (1993). The two different finite element meshes used contain 2200 and 5100 triangular elements, respectively (see Table 2), with a nodal point spacing of approximately 10-15 km. Tests have shown this to be sufficient to resolve the dynamics, since at lower resolution some characteristics of the dynamics are lost whereas a higher resolution results in the same behaviour. The degree of depletion field F was transported using a Lagrangian particle tracer method (300000 to 400000 tracers). Fourth-order Runge-Kutta time integration is used to advect the particle tracers with the mantle flow velocity. The transformation of particle tracer field values (the degree of depletion) to finite element Gauss points uses a Particle in Cell algorithm (Hockney and Eastwood, 1988).

The density effects of the varying composition and phase transitions are accounted for in a buoyancy contribution terms, scaled by the Rayleigh numbers Rb and Rc_k , in the momentum equation (Eq. (3)) (see also De Smet et al., 1998, 2000).

The density of mantle material is related to the degree of depletion F (Jordan, 1979; Vlaar and Van den Berg, 1991). In addition, basalt has its own density and is transformed into heavier eclogite upon reaching depths in excess of 30 km in our model. This depth (corresponding to 0.9 GPa) is somewhat less than the depth of about 40 km (1.2 GPa) which Hacker (1996) states as the minimum pressure of basalt to eclogite transformation. However, rocks formed by partial melting in a hotter mantle are more MgO-rich than present-day MORB (Nisbet, 1982), which may cause the transition to occur at lower pressures and higher temperatures (Green and Ringwood, 1967). Furthermore, garnet may already enter phase assemblages around 0.7-0.8 GPa (Green and Ringwood, 1967; Ito and Kennedy, 1971; Hacker, 1996), raising the bulk density above that of the original basaltic phase assemblage. The kinetics of this transition are approximated assuming a constant relaxation time for the transition of 1.25 My.

We do not consider phase boundary topography due to temperature effects, since we expect this to be dominated by kinetic effects because of the relatively

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

Table 1 Symbols of the energy, momentum, depletion, and rheology (Eqs. (1)–(8))

Symbol	Property	Definition	Value/unit
$\overline{B_1}$	diffusion creep prefactor		Pa s
B_2	dislocation creep prefactor		$Pa^{n_2} s$
C_0	cohesion factor		0 Pa
cp	heat capacity at constant pressure		$1250 \text{ J kg}^{-1} \text{ K}^{-1}$
Di	dissipation number	$\frac{\alpha g h}{c_p}$	
E_1	diffusion creep activation energy	-p	$270 \times 10^3 \text{ J mol}^{-1}$
E_2	dislocation creep activation energy		$485 \times 10^3 \text{ J mol}^{-1}$
e _{ij}	strain rate tensor	$\partial_j u_i + \partial_i u_j$	s^{-1}
е	second invariant of the strain rate tensor	$\left[\frac{1}{2}e_{ii}e_{ii}\right]^{\frac{1}{2}}$	s^{-1}
f(F)	composition dependent viscosity prefactor	$12 - y - y_1$	
F	degree of depletion		
g	acceleration due to gravity		9.8 ms^{-2}
h	length scale		$1005 \times 10^3 \text{ m}$
Н	radiogenic heat production		$W m^{-3}$
n_1	diffusion creep stress exponent		1
n_2	dislocation creep stress exponent		3.25
n _v	yield exponent		10
P	pressure		Ра
R	Gas constant		$8.341 \text{ J} \text{ mol}^{-1} \text{ K}^{-1}$
Ra	thermal Rayleigh number	$\frac{\rho_0 \alpha \Delta T g h^3}{\kappa \eta_0}$	
Rb	compositional Rayleigh number	$\frac{\delta \rho g h^3}{\kappa \eta_0}$	
Rc	phase Rayleigh number	$\frac{\delta \rho g h^3}{\kappa \eta_0}$	
$R_{\rm H}$	internal heating number	$\frac{H_0h^2}{c \kappa \Delta T}$	
S	melt productivity function	c _p r <u></u>	s^{-1}
ΔS	entropy change upon full differentiation		$300 \text{ J kg}^{-1} \text{ K}^{-1}$
t	time		S
Т	temperature		$^{\circ}\mathrm{C}$
T_0	non-dimensional surface temperature		$(273/\Delta T)$
ΔT	temperature scale		2450 °C
V_1	diffusion creep activation volume		$3 \times 10^{-6} \text{ m}^3 \text{ mol}^{-1}$
V_2	dislocation creep activation volume		$17.5 \times 10^{-6} \text{ m}^3 \text{ mol}^{-1}$
W	vertical velocity		ms^{-1}
Ζ	depth		m
ź	unit vector in vertical (downward) direction		
$z_0(T)$	temperature-dependent depth of phase transition		m
α	thermal expansion coefficient	$1 \left(1 - \frac{1}{2} \left(\frac{1}{2} - \frac{1}{2} \left(T \right) \right) \right)$	$3 \times 10^{-5} \text{ K}^{-1}$
Γ	phase function	$\frac{1}{2}\left(1+\sin\left(\pi\frac{z-2o(1)}{\delta z}\right)\right)$	-3
δρ	density difference (chemical or phase)		kg m ⁻³
	basalt		-416 kg m^{-3}
	eclogite		200 kg m^{-3}
	fully depleted peridotite		-226 kg m^{-1}
3.	strain rate		S ¹
ε _y	yield strain rate		10 s '
η	VISCOSITY		Pa s
η_y	yield Viscosity		Pa s $10^{-6} m^2 -1$
к	friction coefficient		10 III S 0.03
μ			0.05

(continued on next page)

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

Table 1 (continued)

Symbol	Property	Definition	Value/unit		
ρ_0	reference density	rı 1 ¹	3416 kg m^{-3}		
τ	second invariant of deviatoric stress tensor	$\left\lfloor \frac{1}{2} \tau_{ij} \tau_{ij} \right\rfloor^2$	Pa		
$\sigma_{\rm n}$	normal stress		Ра		
τ_y	yield stress		Pa		
Φ	viscous dissipation	ηe^2	$J m^{-3} s^{-1}$		

low Clapeyron slope of about 1 MPa/K (Philpotts, 1990).

The phase transitions around 400 and 670 km depth are also taken into account separately for peridotite and eclogite, though without kinetics (i.e., it is assumed to occur instantaneous). The relevant parameters for these phase transitions are given in Table 3. The phase transition of eclogite into

Table 2 Characteristics of the different model experiments

Model	Initial cond	litions	Domain			
	$T_{\rm pot}^{\rm start}$ (°C)	$d_{\rm crust}^{\rm start}$ (km)	No. of elements	No. of boundary points	Width and depth (km)	
M0	1464	0.0	2245	67	1005	
M0b	1464	0.0	2245	67	1200	
M1	1896	20.1	2245	67	1005	
M2	1948	25.6	2245	67	1005	
M3	1976	30.0	2245	67	1005	
M4	2000	35.4	2245	67	1005	
M5	2017	39.8	2245	67	1005	
M6	2051	50.5	2245	67	1005	
Mr	1976	30.0	5101	100	1005	
Ms	1976	29.7	5101	100	1200	
Md	1976	30.0	3395	82 ^a	1005	

Models M0 and M0b are startup runs with increased radiogenic heating (see text). $T_{\rm pot}^{\rm start}$ is the potential temperature and $d_{\rm crust}^{\rm start}$ is the average crustal thickness, both at the start of the experiment. The number of boundary points indicates the number of model points of the triangular elements (of uniform size) used in the discretization of the domain on a single domain boundary and is included as a measure of 1-D resolution. Note that model Md uses a different set of parameters for its phase transitions (peridotite: ~400 km $-\delta\rho/\rho_0=0.0685$, $\gamma=2\times10^6$ Pa K⁻¹; ~670 km $-\delta\rho/\rho_0=0.1552$, $\gamma=-2.5\times10^6$ Pa K⁻¹; basalt: ~400 km $-\delta\rho/\rho_0=0.0392$, $\gamma=2\times10^6$ Pa K⁻¹; ~720 km $-\delta\rho/\rho_0=0.1610$, $\gamma=0.8\times10^6$ Pa K⁻¹) and a heat reservoir at its lower boundary, resulting in a slightly different model evolution.

^a In vertical direction, same resolution as models M1–M6 in lower 804 km of model, and nodal point spacing divided by 2 in horizontal and vertical direction in upper 151 km, with gradual element size change in between. perovskitite lithology in the shallow lower mantle (e.g. Ringwood and Irifune, 1988; Irifune and Ringwood, 1993; Hirose et al., 1999). is not taken into account. Dynamic effects have been predicted of the density inversion between eclogite and postspinel peridotite in the uppermost lower mantle (Irifune and Ringwood, 1993) caused by the deeper position of the transition of eclogite relative to the postspinel transition of peridotite. As recent work has indicated that the transition takes place around 720 km depth (Hirose et al., 1999) rather than around 800 km (e.g. Irifune and Ringwood, 1993), and has a positive Clapeyron slope which would lift the phase transition to even shallower levels in a cool sinking eclogite body, Hirose et al. (1999) predict that 'basaltic crust with perovskitite lithology would gravitationally sink into the deeper mantle' (their p. 53). Earlier numerical experiments of Christensen (1988) show that an eclogite layer trapped in this density inversion zone in the uppermost lower mantle would probably not survive in a convecting mantle. Recent experiments by Kubo et al. (2002), however, indicate that slow kinetics of the lower mantle phase transition of basaltic material may delay the transition on the order of 10 million years, which may increase the residence time and depth extent of the shallow

Table 3								
Parameters	of	the	phase	transitions	used	in	the	models

	-					
Transition	Material	P ₀ (GPa)	Т ₀ (К)	γ (Pa K ⁻¹)	δz (km)	$(\delta \rho / \rho_0)$
400	peridotite	13.4	1756	3×10^{6}	50	0.05
400 670	peridotite	12.3 22.4	1/4/	-2.5×10^{6}	50	0.10

The first column indicates the approximate depth of the transition, P_0 and T_0 are reference pressure and temperature, γ is the Clapeyron slope of the phase transition, δz indicates the depth range over which the transition is smeared out and the last column gives the relative density contrast of the phase transition.

lower mantle density inversion zone. We have validated our assumption of the minor importance of the shallow lower mantle eclogite density inversion by performing an experiment in which it is included (eclogite 430 kg m⁻³ lighter than peridotite between 670 and 720 km depth), which showed no significant stagnation of basaltic material in this density inversion zone.

2.2. Rheology

The deformation of material is accounted for by three separate deformation mechanisms combined in a composite rheological model. The mechanisms included are diffusion creep, dislocation creep, and brittle failure. The first two mechanisms are described by the following expression for the corresponding viscosity component:

$$\eta_i = f(F)B_i \exp\left[\frac{E_i + PV_i}{RT}\right] \tau^{1-n_i}$$
(5)

which is the standard Arrhenius formulation (Karato and Wu, 1993; Van den Berg and Yuen, 1998) with the addition of a composition dependent prefactor f(F). The index *i* indicates the mechanism, either 1 for diffusion creep or 2 for dislocation creep. The other symbols are explained in Table 1. The activation energies E_i and volume V_i are based on Karato and Wu (1993), as is the prefactor B_i , in which an assumed constant and uniform grain size of 1 mm is incorporated. The composition dependent prefactor f(F) has a value of 1 for fertile peridotite and basalt (which have the same viscosity parameters in our models). For depleted peridotite (harzburgite), we apply a prefactor value of 10 for a degree of depletion over 0.05, and a linearly increasing value between F=0.005 and F=0.05 from 1 to 10 (Van Thienen et al., 2003). This is intended to mimic the effect of dehydration during partial melting on the viscosity (Karato, 1986; Hirth and Kohlstedt, 1996; Mei and Kohlstedt, 2000a,b).

For eclogite, the viscosity prefactor f(F) has a value of 0.1. Although Jin et al. (2001) find that for an eclogite with equal amounts of the two main constituents garnet and omphacite, the strength is comparable to harzburgite (which is hydrous in their experiments). Piepenbreier and Stöckhert (2001) find evidence in eclogite microstructures for a much lower

flow strength than predicted from omphacite experimental flow laws. We translate this result to our latter viscosity prefactor value of 0.1.

The third deformation mechanism, brittle failure, is approximated by a yield mechanism which is included to reproduce fracturing-like behaviour when the stress supersedes a certain yield stress τ_y . We use the formulation of Van Hunen et al. (2002)

$$\eta_{\rm y} = \tau_{\rm y} \dot{\boldsymbol{\varepsilon}}_{\rm y}^{-1/n_{\rm y}} \dot{\boldsymbol{\varepsilon}}_{\rm y}^{(1/n_{\rm y})-1} \tag{6}$$

The symbols are explained in Table 1. We prescribe the yield stress τ_y , the yield strain rate ε_y and the yield exponent n_y . The latter describes the brittleness of the behaviour (Van Hunen et al., 2002). We apply a value of 10, which gives a reasonable approximation of pure brittle behaviour. The yield strength τ_y is determined as a function of depth using Byerlee's law (e.g. Lockner, 1995):

$$\tau_{\rm y} = C_0 + \mu \sigma_{\rm n} \tag{7}$$

in which we approximate the normal stress σ_n with the lithostatic pressure (e.g. Moresi and Solomatov, 1998):

$$\tau_{\rm y} = C_0 + \mu \rho g z \tag{8}$$

We use a low value of 0.03 for the friction coefficient μ , consistent with results of Moresi and Solomatov (1998) for the mobilization of the Earth's lithosphere, and assume the cohesion term C_0 to be of minor importance and set it to 0.

The diffusion and dislocation creep viscosities are inversely added to define an effective ductile creep viscosity (Van den Berg et al., 1993) and the minimum of this value and the local yield viscosity is used for the local effective viscosity.

2.3. Fractionation of trace elements

The behaviour of a single incompatible trace element is monitored in the models using the particle tracers. Upon partial melting, the concentration of the trace element is adjusted in each tracer using an equilibrium melting formulation (see e.g. Philpotts, 1990) for each integration time step (which essentially means that we assume equilibration between the melt and the residue during an integration time

step, and consider a new batch of melt in the following time step). As mentioned above, the internal heating rate is a function of trace element concentration in the models. The result is that an enriched crust (generally by a factor 5-20) and a depleted residual mantle are formed. The most important heat producing species in the Archean mantle are U and Th. These have bulk partition coefficients of 1.1×10^{-4} and 1.7×10^{-4} , respectively, for spinel peridotites in equilibrium with a basaltic melt (Beattie, 1993), though temperature, composition, oxygen fugacity and the presence of volatiles may influence these values. We prescribe a distribution coefficient of 10^{-2} for the single incompatible heat producing element in our model which represents both U and Th. The value of the partition coefficient is somewhat larger than the values mentioned above for spinel peridotites. It is however small enough to allow significant fractionation and thus redistribution of heat productivity, but not so small that different degrees of melting result in the same residue concentrations of virtually zero. This allows us to regard the development of our trace element in a more general sense rather than limited to U and Th.

2.4. Model geometry and boundary conditions

For most models, we use a square computational domain of 1005 by 1005 km (see Fig. 1), resulting in an upper mantle aspect ratio of 1.5. One extended model has dimensions of 1200×1200 km, resulting in an upper mantle aspect ratio of 1.8 (see Section 3.4). We use a prescribed velocity on the upper boundary (controlled by basalt generation), periodical side boundaries and free slip conditions on the lower boundary. On the upper boundary, we prescribe a temperature of 0 °C. Although we are dealing with the early Earth, the presence of liquid water on the surface, as indicated by oxygen isotope ratios in zircons of up to 4.4 Gy old (Wilde et al., 2001; Peck et al., 2001), validates this assumption. On the lower boundary, we prescribe a zero heat flux boundary condition to approximate conditions of transient secular cooling of the model by excluding heat input from below.



Fig. 1. Setup of the 2-D square model domain measuring 1005×1005 and 1200×1200 km in respective models listed in Table 2, representing the upper mantle and part of the lower mantle.

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

3. Results

As the initial condition for the models (i.e. the state of the Earth after solidification of the magma ocean) is difficult to reconstruct, we start from a model configuration including an initial basaltic crust and underlying depleted zone. We first test the sensitivity of the long-term development of the models to the initial situation (temperature and initial thickness of crust and depleted zone). The results will be presented in Section 3.1. After this, the dynamics of crustal growth and recycling will be shown from the results of the model calculations (Section 3.2). Next, we will present the long-term development of a number of models in Section 3.3, and finally, the effect of the size of the model lower mantle is considered in Section 3.4.

3.1. Sensitivity to initial conditions

We generate initial situations including a basaltic crust by starting model M0 (and M0b for a model with a larger domain width, see Table 2) with a uniform fertile composition, a geotherm which is below the solidus in the entire domain and a strongly increased internal heating of 250×10^{-12} W/kg. The model rapidly heats up and starts melting, thus producing a crust. Different snapshots from the development of model M0 are used as starting points for a number of cooling and recycling models (M1-M6 in Table 2). Fig. 2 shows the development of the potential temperature and accumulated crustal thickness for the startup model M0. The symbols in Fig. 2 indicate the snapshots from which the cooling and recycling models are started. These are listed, together with their characteristics, in Table 2. In the cooling and recycling models M1–M6, the clock is initially reset to t=0, the internal heating is reset to 15×10^{-12} W/kg (fertile mantle material value representative of the early Earth), and natural decay with a half-life of 2.5 Gy is included.

In the two models with the thickest crust and highest initial temperature (M5 and M6, see Table 2), the entire crust sinks into the mantle within the first few million years of the model evolution. This is a result of the fact that a large part of the crust is transformed into the dense eclogite phase. The upwelling hot mantle material which replaces it locally completely melts, bringing the system back



Fig. 2. Development of the potential temperature and average crustal thickness of the initial model M0. Time progresses from the lower left to the upper right part of the curve. Symbols indicate the snapshots which are used as starting points for different cooling models (see Table 2).

into a magma ocean regime. As our model assumptions are only valid for modest degrees of melting, we do not continue the calculations at complete melting, which is observed here .The results do show that a very thick basaltic crust of 40 km or more thickness is not sustainable on top of a hot, weak mantle in the early Earth.

The four somewhat cooler models M1–M4 (see Table 2) show higher rates of activity (vigor of convection and melt productivity) for higher initial temperature and greater crustal thickness. As a result of the higher crustal production rate, a higher cooling rate is observed for the hotter models, so that their volume averaged temperatures and surface heat fluxes converge after some hundreds of millions of years.

3.2. Dynamics of crustal growth and recycling

Three distinctive types of dynamics can be discerned in the process of crustal growth and recycling. The first is a small-scale circulation near the bottom of the crust which transports eclogite down into the mantle and which produces new crust by partial melting of the upwelling part of the circulation. Fig. 3 shows the development of such a coupled system in model Mr. The left frames show the composition field (individual particle tracers are plotted for basalt (black) and eclogite (red)), the

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



10

middle frames the temperature field (colours) and circular to oblong melt production zones (blue contours), and the right-hand side frames show the viscosity field with 10%, 50% and 90% isolines (dashed-solid-dashed) for the phase transitions of peridotitic (black lines) and basaltic (white lines) material listed in Table 3. In Fig. 3a, no eclogite delamination is taking place yet. Two regions of thickened crust are visible around x=500 km and x=750 km, and partial melting occurs in the upwelling flow near the right-hand side region. This is illustrated by the melting contours in the temperature frame. The melt which is produced is instantaneously transported vertically to the surface where it adds new material to the locally thickened crust. In the next frame (Fig. 3b), eclogite starts to move downwards. At the same time, we see a similar but mirrored circulation arising below the left-hand side thickened crustal region. About 4 million years later (Fig. 3c), active delamination can be observed in both circulation systems, with high velocities in the common downwelling limb of the shallow circulations, and stretches of eclogite and depleted peridotite piling up in the lowermost upper mantle. Partial melting continues in the upwelling parts as indicated by the blue contours in the corresponding temperature snapshot. The recycling of eclogite into the mantle continues for tens of millions of years (see Fig. 3d, 29 million years later).

Larger-scale upwellings may add significantly more material to the crust. Fig. 4 illustrates the development of a diapir originating from a lower mantle upwelling. As in Fig. 3, the left-hand frames show the composition field, the middle frames the temperature field and the melting zone (which is quite large in this case), and the right-hand side frames the viscosity field with phase transitions. Because of its lower mantle origin, the material is both hot and fertile, indicated by the deep blue colour, and starts melting at 400 km depth (which is the imposed maximum depth of melting in the model) in Fig. 4a. Melting continues up to about 150 km depth, where the diapiric ascent is stopped by the rapid increase of the temperature dependent viscosity (Fig. 4b, righthand side frames; note that the identical times of snapshots a and b indicate extremely fast upwelling in less than 10 ky). The diapir head spreads out and the left-hand side rises further (Fig. 4c) and melts up to a degree of nearly 0.30. The original diapir spreads out over nearly the entire domain of 1005 km. The large volume of melt produced locally thickens the crust by over 40 km (Fig. 4d). The time scale on which this event takes place is extremely short. Although inflow of lower mantle material and partial melting continue for some millions of years through the narrow lowviscosity plume tail, the rise of the diapir head through the upper mantle takes place in less than 20 ky. This is the effect of the nonlinear component of the rheology, as has been shown in the literature (Weinberg and Podladchikov, 1994; Larsen et al., 1997; Larsen and Yuen, 1997; Van Keken, 1997; Van Thienen et al., 2003), and is illustrated by the very low viscosities down to about 10^{17} Pa s in the plume head and tail. As is shown by the composition snapshots of Figs. 4d and 5a, less than half of the original lower mantle material enters the upper mantle through this event.

At least as catastrophic is the third type of dynamics, which is characterized by large-scale delamination of a thickened crust into the mantle. Fig. 5 shows the entire domain, starting just after the diapir event shown in Fig. 4. The deeper part of the crust which is significantly thickened turns into eclogite in about 2 million years (Fig. 5a) and sinks into the mantle. This is facilitated by a weakening of the crust by the yielding mechanism, resulting in a low effective viscosity in the upper part of the sinking material in the upper right of the viscosity frame of Fig. 5a. The middle frame of Fig. 5a shows that partial melting is taking place in the zone where material is sucked into the zone of upwelling created by the sinking crustal block. In this region, high flow velocities cause a significant lowering of the viscosity through the nonlinear component of the viscosity. As

Fig. 3. Four snapshots of composition, temperature, and viscosity from model Mr show how small-scale circulations near the base of the crust allow recycling of eclogite into the mantle and production of new crust by partial melting of upwelling mantle peridotite. In the left-hand side panels, the colour scale from blue to orange indicates the degree of depletion F. Black and red tracers represent basalt and eclogite, respectively. The middle panels show the temperature field (colour scale in centigrade) and areas were partial melting is taking place (inside the contours). The right-hand side panels show the viscosity field, and the location of phase transitions for peridotitic (black) and basaltic (white) compositions, at 50% (solid) and 10% and 90% (dashed) of the transition.

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 4. Four snapshots of composition, temperature, and viscosity from model Mr show the ascent and decompression melting of a mantle diapir, originating from the lower mantle. For an explanation of the colours, see the caption of Fig. 3.

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 5. Four snapshots of composition, temperature, and viscosity from model Mr show the sinking of the entire crust into the mantle, driven by a large mass of dense eclogite. For an explanation of the colours, see the caption of Fig. 3.

the crustal block is quite cold and strong (illustrated by the viscosity snapshots), it pulls down with it all the existing crustal material, which is mostly in the basalt phase and positively buoyant, see Fig. 5b. This process is essentially comparable to the lithospheric collapse modelled by Moresi and Solomatov (1998), but in our model, chemical buoyancy is an important factor in causing the collapse. Melting continues in the upper mantle, creating a new basaltic crust (Fig. 5c). Note that this crust is significantly thinner (about 30 km) than the crust shown in Fig. 5b. After the downgoing crust is sufficiently thinned, it breaks off at the weak zone near the surface (see Fig. 5c, viscosity panel), caused by the yielding mechanism, and further 'subduction' is halted. The basalt which was pulled down into the mantle has completely transformed into eclogite by now and the entire 'crustal lobe' sinks into the deeper mantle (Fig. 5d). The complete replacement of the old crust by a new crust is illustrated by Fig. 6, showing the variations of the formation age of crustal material. The figure, corresponding to the snapshot shown in Fig. 5d, shows the old crust (blue) as it is sinking into the mantle, and the freshly produced crust (green, yellow, orange, purple) forming the new top of the domain. Note that the old crust has pulled down a significant amount of young crust (purple). In the upper part of the lower mantle, the subducted crust slowly heats up and becomes weaker. It breaks up allowing lower mantle peridotite to break into the upper mantle, while sinking crust replaces this material in the lower mantle (Fig. 5d). Note the very short time scale of the whole process, where the replacement of the entire shallow crust takes place in less than 2 million years. The settling of the crust into the lower mantle to the bottom of the model domain takes much longer, on the order of 300 million years as both the density contrast between the sinking crust and the mantle decrease and the viscosity increases with depth (and decreasing strain rate).

In model Mr, lower mantle diapirism triggers the resurfacing event by adding large amounts of basalt to the crust, which becomes unstable. However, these two types of dynamics are not necessarily connected and at least resurfacing can exist on its own. This is illustrated in Fig. 7, where a resurfacing event takes place which is triggered by a crust which is thickened not by lower mantle diapirism but by partial melting in the convecting upper mantle in model Md (see Table 2). Diapirism which does not trigger a resurfacing event is not often observed in our model results, since diapirs generally add enough material to a crustal section to make in unstable.

3.3. Long-term thermal and chemical development

The thermal evolution of model Mr is illustrated in Fig. 8a by the solid black curve, showing the volume averaged temperature as a function of time. The dashed curves show the volume-averaged temperatures of the lower (670-1005 km depth, long dashes) and upper (150-670 km depth, short dashes) mantle. Three stages can be discerned. During the first stage, lasting about 230 My, small-scale delamination and crustal growth (see Fig. 3) generate significant cooling of at a rate of about 300 K/Gy (and crustal growth, as shown in Fig. 8b), mainly in the upper mantle. The second, quite short, stage is characterized by rapid crustal growth and associated cooling, caused by lower mantle diapir ascent and complete resurfacing as shown in Figs. 4 and 5. A drop in the volumeaveraged temperature of about 200 °C can be observed in the complete domain, with a much stronger drop for the lower mantle. An average (over the entire domain width) of over 40 km of crust is produced. The third stage shows an absence of crustal growth, and because of this and the fact that the lower mantle temperature results in a thicker lithosphere, the cooling rate for the complete domain is reduced relative to the first two stages. The lower mantle slowly heats up again. Fig. 8c shows the root mean square velocity for the entire domain as a function of time. The three stages observed in the temperature and crustal production time series can also be seen here. During the first stage, strong variations over short periods of time are observed, with rms velocities ranging from about 1 cm/year to about 10 m/year. The stage of rapid crustal growth is characterized by extreme rms velocities of about 10^2 m/year. In the subsequent quiet stage, the rms velocity varies between roughly 3 mm/year and 10 cm/year, on significantly longer time scales than in the first stage. No absolute numbers for cooling rates are given here since the small model domain (1005 km depth) does not allow comparison to the Earth, as will be discussed below (see Section 3.4, where the larger

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 6. The formation age of every basaltic/eclogitic tracer in the domain of model Mr at 237.32 My is shown in colour. Tracer ages outside the colour scale are dark blue (older) or green (younger). The complete domain of 1005×1005 km is shown.

model domain width and depth model Ms will be discussed).

The evolution of the incompatible trace element distribution of model Mr is illustrated in Fig. 9, which shows the incompatible trace element concentration at different times during its evolution. The initial concentration of the trace element is 1, and the distribution coefficient is 10^{-2} . The 10 frames in this figure show the concentration field at 10 subsequent times during the evolution of the model. Initially, the lower mantle is pristine. The upper mantle shows an increasing trace element depletion for shallower depths, where the mantle peridotite is subject to melting. This shallow partial melting also produces the enriched basaltic layer at the top boundary. After some 15 million years, slightly depleted upper mantle material is entrained into the lower mantle flow. As the recycling of eclogite into the mantle progresses, we see an increasing amount of grey-coloured enriched eclogite 'blobs' in the upper mantle (frames at 154.6 and 228.9 My). The progressive cooling and the accumulation of eclogite blobs in the deepest upper mantle forces a breakdown of completely separate lower mantle and upper mantle convection (frames at 230.4-241.91 My), as predicted by Christensen and Yuen (1985). This brings more highly depleted material and enriched eclogite blobs into the lower mantle. Over the following 300 million years, the enriched crust sinks deeper into the lower mantle and settles at the bottom boundary. The upper and lower mantle continue to mix and form a more or less uniform reservoir of moderately depleted peridotite in which blobs of eclogite are dispersed. The high content of heat producing elements of the settled crust in the lower mantle makes it a heat source for convection in the mantle above, which may entrain some material from this layer. But essentially it remains stable for the remaining 500 million years of the model run.

The development of the bulk and incompatible trace element chemistry of this model (Mr) is presented in Fig. 10. Time series of the averaged degree of depletion of the peridotite present are shown in Fig. 10a for the entire domain, and for the lithosphere (0–150 km depth), the upper mantle (150–670 km depth) and the lower mantle (670–1005 km depth) separately. During the first 230 million years, we see a steady increase in the average degree of depletion of the entire domain of model Mr (Fig. 10a). The lower mantle stays nearly pristine



P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 7. Three snapshots of model Md, showing that diapirism and resurfacing, which take place nearly simultaneously in model Mr (see Figs. 4 and 5) are not necessarily connected. Model Md shows a crustal section which has thickened without material addition by a diapir triggering a resurfacing event.

during this entire period. The upper mantle doubles its average degree of depletion over this period of time. However, as the interaction between the upper and lower mantle increases due to the cooling of the model and the accumulation of eclogite blobs at the bottom of the upper mantle, some of the depleted material of the upper mantle is dumped into the lower mantle. The lithospheric degree of depletion fluctuates but

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 8. Time series of (a) volume averaged temperature for the entire model domain (solid line), the lower mantle (670–1005/1200 km depth, long dashes) and the upper mantle (150–670 km depth, short dashes), (b) accumulated amount of created crust represented as an average thickness over the entire domain with, (c) and root mean square velocity, for models Mr and Ms (larger domain of 1200 km depth).

shows a general decrease because more material is mixed into the upper mantle. After this initial stage, two important events (intrusion of a lower mantle diapir into the upper mantle and subsequent recycling of the thickened crust into the mantle) cause a marked increase in the averaged degree of depletion of the domain. The increase is observable in every reservoir, and practically at the same time (230 My). After this, the slow sinking of the crust and associated depleted mantle from the upper into the lower mantle causes an increase in the lower mantle value and a simultaneous decrease in the upper mantle value. Continued exchange of material between the upper and lower mantle reduces the difference over the following hundreds of million years, in line with the effective mixing illustrated in Fig. 9.

Fig. 10b shows the development of the approximate volume-averaged incompatible trace element concentrations in the lithosphere, upper mantle and lower mantle. The same distinction in two separate phases can be made as in Fig. 10a. During the first 230 million years, a steady increase in the average incompatible isotope concentration can be observed in the lithosphere. This is due to the accumulation of crust with a high concentration of the incompatible element. Complementary to this trend (but less pronounced because the reservoir is larger), we see a decrease in the average incompatible element concentration of the upper mantle, since the depleted residue which is formed upon partial melting is recycled through the upper mantle. The lower mantle shows a slight decrease in the concentration, which is due to radioactive decay. In the other curves, this is also present but invisible due to the larger effect of differentiation.

The two large-scale events around 230 My (lower mantle diapir and large-scale sinking of the crust into the mantle) have a strong impact on the average incompatible trace element distribution. As the crust forms an important reservoir for this element, its sinking into the lower mantle causes a strong decrease in the average trace element concentration in the lithospheric zone. Two causes can be named. Firstly, the new crust which is formed is much less voluminous than the old crust which has been subducted. And secondly, the new crust was formed from already depleted mantle, so it will have lower average incompatible trace element concentrations. After these events, some entrainment of eclogite tracers from the settled crust at the bottom boundary of the model into the upper mantle somewhat increases the upper mantle average concentration and decreases it in the lower mantle. The effect of radioactive decay is visible in each curve.

3.4. Lower mantle size

As has been noted before, the lower mantle in our models is significantly smaller than in the Earth. Including a larger lower mantle at the same resolution,



Fig. 9. Ten snapshots of the trace element concentration field illustrate the chemical differentiation of model Mr. Pristine, undifferentiated material is white. Material which is depleted in trace elements is yellow, and the enriched melt products are blue. The distribution coefficient applied in 10^{-2} .

P. van

Thienen et al.

<

Tectonophysics xx (2004) xxx-xxx

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx



Fig. 10. The chemical evolution of model Mr is presented in the form of time series for the averaged degree of depletion (a) and the averaged trace element concentration (b). Averaged values for the entire domain are shown in black. A subdivision has been made into the lithosphere (0–150 km depth, red curves), the upper mantle (150–670 km depth, green curves) and the lower mantle (670–1005 km depth, blue curves). Colours refer to the version on the web.

however, is computationally quite expensive. The size of the lower mantle is important for a number of reasons. It is a reservoir for both heat and undepleted peridotite. The models as presented in the previous sections are therefore capable of producing less crust than the Earth and they show a more rapid cooling than expected for the Earth.

Another important aspect is the layering of mantel convection. Christensen and Yuen (1985) showed that the layering of mantle convection is a function of the Rayleigh number (see Table 1 for symbol definitions):

$$Ra = \frac{\rho_0 g \alpha \Delta T h^3}{\eta_0 \kappa} \tag{9}$$

In general, high Rayleigh number systems tend to develop a layered convection pattern in the presence

of an endothermic phase boundary, whereas low Rayleigh number systems would tend to a whole mantle convection pattern. This suggests that the breakdown of layered convection as takes place, e.g. model Mr around 230 My (see Figs. 4 and 5) would be delayed in a system with a higher Rayleigh number. In order to test the sensitivity of the results to the depth of the lower mantle part of the model, we have done experiments with an increased domain width (and depth) of 1200 km (model Ms, see Table 2), resulting in a lower mantle depth of 530 km. The resulting dynamics of the extended model described above are summarized in the time series of volumeaveraged temperature, total amount of crust produced, and root mean square velocity of the entire domain, in Fig. 8. Curves for the base model Mr can be used for comparison. As is illustrated by both strong drops in temperature and strong increases in the amount of crust produced, model Ms has a similar resurfacing event as described in Section 3.2 and shown in Fig. 5 for model Mr. The model shows a higher melt productivity than the smaller model Mr at a comparable cooling rate. This is a direct result of the reduced surface to volume ratio. Furthermore, the higher volume of fertile peridotite which can act as a source rock for basaltic material allows more crust to be produced. The extended model Ms also shows continued melting (although at a low rate) after the resurfacing event, in contrast to the behaviour of model Mr, which shows no more melting after resurfacing.

Although because of the higher Rayleigh number of model Ms, it would be expected that layered convection is maintained for a longer period of time in this models than in model Mr (only 1005 km depth), the extended model shows a somewhat higher crustal production rate and generate a thick unstable crust on a shorter time scale, which causes resurfacing on a shorter time scale as well. Clearly, the fact that we are dealing with thermo-chemical convection rather than merely thermal convection frustrates predictions based on the results of Christensen and Yuen (1985) and Steinbach et al. (1993) who find stronger layered convection for higher Rayleigh numbers. For the Earth, episodic switching between whole mantle convection and layered mantle convection has been inferred from geochemistry (Stein and Hofmann, 1994) and modelled by Breuer and Spohn (1995). In

the extended model Ms, partially layered convection does take place after the resurfacing events, but a complete separation of upper and lower mantle is no longer observed.

4. Discussion

4.1. Large Archean mantle plumes

The strong temperature dependence of the viscosity of the mantle causes the maximum horizontal temperature variations which can be created in the convecting mantle to be a function of mantle temperature (McKenzie and Bickle, 1988). For the present mantle, plume excess temperatures are estimated to be up to 200 to 250 K (Herzberg and O'Hara, 1998). In a hotter Archean mantle, this would be reduced to about 50 to 150 K (Nisbet et al., 1993). However, the mantle diapir which enters the upper mantle in model Mr (Fig. 4) shows excessive melting due to its high excess temperature of about 250 K. Since this excess temperature is directly related to the stability of the phase transition around 670 km, and its breakthrough in Fig. 4 occurs on a much smaller scale than the width of the domain, this absolute excess temperature has significance despite the limited extent of the lower mantle and a model aspect ratio which is quite different from that of the Earth. Clearly, the argument of reduced maximum excess temperatures for plumes in a hotter mantle of McKenzie and Bickle (1988) and Nisbet et al. (1993) breaks down for plumes originating from a breakthrough between layered convection cells as illustrated here. This demonstrates that in a hotter mantle, which will show a stronger layering of mantle convection (Christensen and Yuen, 1985), plumes with high excess temperatures and massive melting are feasible.

4.2. Rapid resurfacing

In Section 3.2, we have seen that the entire crust may periodically sink into the mantle (see Fig. 5). In the case of such an event, it is completely replaced by new crust, and a large amount of heat is released from the mantle. The crust sinks because of gravitational instability, caused by the transformation of basalt into eclogite in a crust significantly thickened by the breakthrough of lower mantle material into the upper mantle. An important parameter is the rheology of the sinking crust. In Fig. 5, it is strong enough to pull down the entire existing (and partly positively buoyant) crust, such that a strong downward force localized at the site of the thickened crust causes resurfacing of the entire model. This type of dynamics resembles models for the dynamics of Venus during its history. Crater counts on the surface of Venus indicate that this planet underwent a global resurfacing event about 500 million years ago (Schaber et al., 1992). Different mechanisms have been proposed to explain the resurfacing event(s). Herrick and Parmentier (1994) suggest on the basis of thermal evolution calculations that the episodic reversal of two layers in the mantle may trigger a period of massive volcanism which completely renews the crust. Solomatov and Moresi (1996) suggest that plate tectonics were active on Venus before 500 My, which stopped because the stresses in the lithosphere dropped below the yield strength. A similar argument is used by Fowler and O'Brien (1996) to explain episodic heating of the mantle under a thickening stagnant lid, which experiences increasing buoyancy induced stresses during its cooling up to the point where plastic failure occurs and the lithosphere can subduct into the mantle. Turcotte (1995) compares 'episodic subduction' (p. 16935) on Venus to the foundering of the solidified crust on a cooling lava lake. Due to the cooling of the crust, it becomes gravitationally unstable and sinks into the lake in parts which are separated by cracks. Our results most resemble model of Parmentier and Hess (1992), in which the lithosphere episodically becomes negatively buoyant due to cooling, causing it to sink into the mantle. Secondly, the basalt to eclogite phase transition in the lower crust causes parts of it to become negatively buoyant and recycle into the mantle. The material sinking into the mantle makes room for new partial melting of mantle peridotite, generating new crustal material and a depleted root, which can start to cool again. The results of Parmentier and Hess (1992) indicate that this process could take place on Venus with a periodicity of 300 to 500 million years. No spatial scale of recycling of lithosphere into the mantle is included in their model. Our results show recycling of eclogite and also depleted peridotite into the mantle on two scales:

small-scale delaminations and large-scale sinking of the entire lithosphere into the mantle. However, Parmentier and Hess (1992) assume that eclogite which sinks into the mantle completely mixes with it, also chemically, essentially refertilizing the mantle. The mechanism of mantle refertilization by reaction with eclogite (Yaxley and Green, 1998) is not included in our model. Furthermore, as our model domain includes only a small part of the lower mantle corresponding to the width of the domain, a reduced amount of fertile peridotite and heat is present in our models, which further limits the maximum amount of crust which may be produced.

4.3. Thermal evolution of the mantle

Although the limited size of the lower mantle and the geometry and aspect ratio of our models does not allow quantitative statements on the absolute cooling rate of the Earth, the results do show that specifically the resurfacing mechanism as illustrated in Fig. 5 is a strong cooling agent. The volume-averaged temperature curves of Fig. 8 show temperature decrease values of about 100 to 200 $^{\circ}$ C for a single resurfacing event in each model. We speculate that such a mechanism may have caused rapid cooling in the early history of the Earth over a period of several hundreds of millions of years until the temperature dropped to levels where the mechanisms described in this work are no longer viable.

4.4. Geochemical evolution of the mantle

Geochemical studies indicate that the earliest rocks found on Earth were already derived from a depleted mantle (Hamilton et al., 1983; Patchett, 1983; Vervoort et al., 1996; Blichert-Toft et al., 1999). Based on Hf isotope data, this depleted mantle was derived from a uniform chondritic reservoir at least 4.08 billion years before present (Amelin et al., 2000). Furthermore, the compiled isotope data from different Archean terranes suggest that during the Archean the mantle source of greenstone belts became more depleted with time (Condie, 1990; Amelin et al., 2000). Our model results are consistent with this observation. Fig. 10 shows a steady, though not constant, increase of the averaged extracted melt degree for the upper mantle during the first 230 million years of the model evolution for model Mr. The same trend is illustrated in the incompatible element concentration plots for this model in Fig. 9, which shows increasingly darker colors (i.e. lower concentrations) in the model upper mantle during the first 230 million years. The small extent of this model lower mantle limits the amount of fertile mantle peridotite which may be transported into the upper mantle. Influx of fresh lower mantle material would lower the average degree of depletion in the upper mantle, as can be observed in Fig. 10a between 230 and 300 million years.

The range in trace element compositions found in basalts from different geodynamical settings on the present Earth indicates that the mantle contains several more or less separate geochemical reservoirs (Hofmann, 1997). Tackley (2000) gives an overview of different models which have been suggested to explain the presence of these reservoirs in the mantle. Furthermore, trace element balance calculations require a 'hidden' reservoir of enriched material apart from the continental crust and the depleted mantle reservoirs to comply with the assumption of a chondritic bulk silicate Earth (e.g. Galer and Goldstein, 1991; McDonough and Sun, 1995). This is also supported by the observation of subchondritic ratios of Nb/La, Nb/Ta, and possibly Ti/Zr in continental crust, depleted mantle and mid-ocean ridge basalt (Rudnick et al., 2000). Rudnick et al. (2000) propose that the missing Nb, Ta and possibly Zr reside in a reservoir of eclogite that underwent partial melting during subduction. They calculate that this reservoir should have a mass of about 0.5% to 6% of the silicate Earth to compensate for the deficit. The subduction and storage of eclogite in the lower mantle as a mechanism of creating an enriched reservoir in the lower mantle has been investigated in numerical convection models by Christensen and Hofmann (1994). Our model results show that subduction in the plate tectonics sense is not required to bring large amounts of eclogite into the lower mantle, and keep it there. Though the small-scale delamination as described in Section 3.2 and shown in Fig. 3 only generates more or less uniformly distributed blobs of eclogite in the entire model mantle, large-scale crustal sinking as shown in Fig. 5 does generate an enriched reservoir (see Fig. 9) in the lower mantle which is stable for hundreds of millions of years at least. In

recent years, geodynamical models have been proposed featuring a dense enriched layer in the bottom third of the mantle (Coltice and Ricard, 1999; Kellogg et al., 1999; Van der Hilst and Kárason, 1999; Albarède and Van der Hilst, 2002). We speculate on the basis of our modelling results that such a layer may have been formed during a short time window in the early evolution of the Earth's mantle by compositional differentiation in pressure release partial melting of mantle peridotite and subsequent large scale sinking of the thick basaltic/eclogitic crust into the lower mantle as an alternative to or precursor of the subduction model proposed by Albarède and Van der Hilst (2002).

5. Conclusions

As argued in Introduction, the geodynamics of the early Earth were probably very different from the recent situation. Our model results, relevant for the hotter conditions in the early Earth, indicate that a number of processes may have been important:

- Small-scale $(x \times 10^2 \text{ km})$ circulations of the shallow upper mantle and the lower crust, with partial melting in the upwelling limb of the convection and delamination and recycling of eclogitic lower crust in the downwelling limb, active over extended periods of time $(x \times 10^6 \text{ year})$.
- Episodic large-scale resurfacing events, in which a large part of or even the entire crust sinks into the mantle and is replaced by new crust at the surface. These events are quite rapid and take place within a few millions of years.
- Very fast penetration (5 m/year) of lower mantle diapirs into the upper mantle, bringing hot and fertile mantle material to supersolidus conditions. This process generates large amounts of melt which may locally significantly thicken the crust (x×10 km) and subsequently trigger the aforementioned process of resurfacing.

An important consequence of the latter process is that the penetration of lower mantle diapirs into the upper mantle in a layered convection system allows plume excess temperatures which are significantly higher (about 250 K) than the maximum plume excess temperatures which are predicted from the modified boundary layer theory of McKenzie and Bickle (1988) as applied to the Archean (about 50–150 K) by Nisbet et al. (1993). The episodic resurfacing in particular may have contributed to the formation of a bottom reservoir enriched in incompatible elements, as the 'subducted' crust sinks into the model lower mantle and forms a stable layer at the bottom boundary. We speculate that such a process may be responsible for the complete resurfacing of Venus which is dated at about 500 My from crater counts.

Acknowledgments

We would like to thank Tine Larsen and an anonymous reviewer for constructive reviews, which helped improve the manuscript. This work was sponsored by the Stichting Nationale Computerfaciliteiten (National Computing Facilities Foundation, NCF) for the use of supercomputer facilities, with financial support from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (Netherlands Organization for Scientific Research, NWO). Further computational facilities used were funded by ISES (Netherlands Research Centre for Integrated Solid Earth Science) which is gratefully acknowledged.

References

- Abe, Y., 1993. Thermal evolution and chemical differentiation of the terrestrial magma ocean. In: Takahashi, E., Jeanloz, R., Rubie, D. (Eds.), Evolution of the Earth and Planets. Geophysical Monograph, vol. 74. AGU, Washington, DC, pp. 41–54.
- Abe, Y., 1997. Thermal and chemical evolution of the terrestrial magma ocean. Phys. Earth Planet. Inter. 100, 27–39.
- Albarède, F., Van der Hilst, R.D., 2002. Zoned mantle convection. Philos. Trans. R. Soc., A 360 (1800), 2569–2592.
- Amelin, Y., Lee, D.-C., Halliday, A.N., 2000. Early-middle Archaean crustal evolution deduced from Lu–Hf and U–Pb isotopic studies of single zircon grains. Geochim. Cosmochim. Acta 64 (24), 4205–4225.
- Beattie, P., 1993. The generation of uranium series disequilibria by partial melting of spinel peridotite: constraints from partitioning studies. Earth Planet. Sci. Lett. 117, 379–391.
- Blichert-Toft, J., Albarède, F., Rosing, M., Frei, R., Bridgwater, D., 1999. The Nd and Hf isotopic evolution of the mantle through the Archean. Results from the Isua supracrustals, West Greenland, and from the Birimian terranes of West Africa. Geochim. Cosmochim. Acta 63 (22), 3901–3914.

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

- Breuer, D., Spohn, T., 1995. Possible flush instability in mantle convection at the Archaea–Proterozoic transition. Nature 378, 608–610.
- Christensen, U.R., 1988. Is subducted lithosphere trapped at the 670-km discontinuity? Nature 336, 462–463.
- Christensen, U.R., Hofmann, A.W., 1994. Segregation of subducted oceanic crust in the convecting mantle. J. Geophys. Res. 99 (B10), 19867–19884.
- Christensen, U.R., Yuen, D.A., 1985. Layered convection induced by phase transitions. J. Geophys. Res. 99, 10291–10300.
- Coltice, N., Ricard, Y., 1999. Geochemical observations and one layer mantle convection. Earth Planet. Sci. Lett. 174, 125–137.
- Condie, K.C., 1990. Geochemical characteristics of Precambrian basaltic greenstones. In: Hall, R.P., Hughes, D.J. (Eds.), Early Precambrian Basic Magmatism. Blackie, Glasgow, pp. 40–55.
- De Smet, J.H., 1999. Evolution of the continental upper mantle: numerical modelling of thermo-chemical convection including partial melting. PhD thesis, Utrecht University, The Netherlands.
- De Smet, J.H., Van den Berg, A.P., Vlaar, N.J., 1998. Stability and growth of continental shields in mantle convection models including recurrent melt production. Tectonophysics 296, 15–29.
- De Smet, J.H., Van den Berg, A.P., Vlaar, N.J., 2000. Early formation and long-term stability of continents resulting from decompression melting in a convecting mantle. Tectonophysics 322, 19–33.
- Dupeyrat, L., Sotin, C., 1995. The effect of the transformation of basalt to eclogite on the internal dynamics of Venus. Planet. Space Sci. 43 (7), 909–921.
- Fowler, A.C., O'Brien, S.B.G., 1996. A mechanism for episodic subduction on Venus. J. Geophys. Res. 101 (E2), 4755–4763.
- Galer, S., Goldstein, S., 1991. Early mantle differentiation and its thermal consequences. Geochim. Cosmochim. Acta 55, 227–239.
- Green, D.H., Ringwood, A.E., 1967. An experimental investigation of the gabbro to eclogite transformation and its petrological applications. Geochim. Cosmochim. Acta 31, 767–833.
- Hacker, B.R., 1996. Eclogite formation and the rheology, buoyancy, seismicity and H₂O content of oceanic crust. Subduction: Top to Bottom, AGU Monogr., pp. 337–346.
- Hamilton, W.B., 1998. Archean magmatism and deformation were not products of plate tectonics. Precambrian Res. 91, 143–179.
- Hamilton, P.J., O'Nions, R.K., Bridgwater, D., Nutman, A., 1983. Sm–Nd studies of Archaean metasediments and metavolcanics from West Greenland and their implications for the Earth's early history. Earth Planet. Sci. Lett. 62, 263–272.
- Helmstaedt, H.H., Scott, D.J., 1992. The Proterozoic ophiolite problem. In: Condie, K.C. (Ed.), Proterozoic Crustal Evolution. Elsevier, Amsterdam, The Netherlands, pp. 55–95.
- Herrick, D.L., Parmentier, E.L., 1994. Episodic large-scale overturn of two-layer mantles in terrestrial planets. J. Geophys. Res. 99 (E1), 2053–2062.
- Herzberg, C., O'Hara, M.J., 1998. Phase equilibrium constraints on the origin of basalts, picrites and komatiites. Earth-Sci. Rev. 44 (1–2), 39–79.

- Herzberg, C., Zhang, J., 1996. Melting experiments on anhydrous peridotite KLB-1: compositions of magmas in the upper mantle and the transition zone. J. Geophys. Res. 101, 8271–8295.
- Hirose, K., Fei, Y., Ma, Y., Mao, H.-K., 1999. The fate of subducted basaltic crust in the earth's lower mantle. Nature 397, 53–56.
- Hirth, G., Kohlstedt, D.L., 1996. Water in the oceanic upper mantle: implications for rheology, melt extraction and the evolution of the lithosphere. Earth Planet. Sci. Lett. 144, 93–108.
- Hockney, R.W., Eastwood, J.W., 1988. Computer Simulations Using Particles. IOP Publishing, Bristol.
- Hofmann, A.W., 1997. Mantle geochemistry: the message from oceanic volcanism. Nature 385, 219–229.
- Irifune, T., Ringwood, A.E., 1993. Phase transformations in subducted oceanic crust and buoyancy relations at depths of 600–800 km in the mantle. Earth Planet. Sci. Lett. 117, 101–110.
- Ito, K., Kennedy, G.C., 1971. An experimental study of the basalt– garnet granulite–eclogite transition. In: Heacock, J.G. (Ed.), The Structure and Physical Properties of the Earth's Crust. American Geophysical Union, Washington, DC, pp. 303–314.
- Jaques, A.L., Green, D.H., 1980. Anhydrous melting of peridotite at 0–15 kb pressure and the genesis of tholeiitic basalts. Contrib. Mineral. Petrol. 73, 287–310.
- Jin, Z.-M., Zhang, J., Green II, H.W., Jin, S., 2001. Eclogite rheology: implications for subducted lithosphere. Geology 29 (8), 667–670.
- Jordan, T.H., 1979. Mineralogies, densities, and seismic velocities of garnet lherzolites and their geophysical implications. The Mantle Sample: Inclusions in Kimberlites and Other Volcanics. American Geophysical Union, Washington, DC, pp. 1–14.
- Kameyama, M., Fujimoto, H., Ogawa, M., 1996. A thermochemical regime in the upper mantle in the early earth inferred from a numerical model of magma-migration in a convecting upper mantle. Phys. Earth Planet. Inter. 94, 187–215.
- Karato, S., 1986. Does partial melting reduce the creep strength of the upper mantle? Nature 319, 309–310.
- Karato, S.I., Wu, P., 1993. Rheology of the upper mantle: a synthesis. Science 260, 771–778.
- Kellogg, L., Hager, B.H., Van der Hilst, R.D., 1999. Compositional stratification in the deep mantle. Science 283, 1881–1884.
- Kubo, T., Ohtani, E., Kondo, T., Takumi, K., Toma, M., Hosoya, T., Sano, A., Kikegawa, T., Nagase, T., 2002. Metastable garnet in oceanic crust at the top of the lower mantle. Nature 420, 803–806.
- Kusky, T.M., Li, J.-H., Tucker, R.D., 2001. The Archean Dongwanzi ophiolite complex, North China craton: 2.505-Billionyear-old oceanic crust and mantle. Science 292, 1142–1145.
- Larsen, T.B., Yuen, D.A., 1997. Ultrafast upwelling bursting through the upper mantle. Earth Planet. Sci. Lett. 146, 393–399.
- Larsen, T.B., Yuen, D.A., Smedsmo, J.L., Malevsky, A.V., 1997. Generation of fast timescale phenomena in thermo-mechanical processes. Phys. Earth Planet. Inter. 102, 213–222.
- Lockner, D.A., 1995. Rock failure. In: Ahrens, T.J. (Ed.), Rock Physics and Phase Relations. A Handbook of Physical Constants. American Geophysical Union, Washington, DC, pp. 127–147.

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

- McCulloch, M.T., Bennett, V.C., 1994. Progressive growth of the Earth's continental crust and depleted mantle: geochemical constraints. Geochim. Cosmochim. Acta 58 (21), 4717–4738.
- McDonough, W.F., Sun, S.-s., 1995. The composition of the Earth. Chem. Geol. 120, 223–253.
- McFarlane, E.A., Drake, M.J., Rubie, D.C., 1994. Element partitioning between Mg-perovskite, magnesiowüstite, and silicate melt at conditions of the Earth's mantle. Geochim. Cosmochim. Acta 58 (23), 5161–5172.
- McKenzie, D.P., Bickle, M.J., 1988. The volume and composition of melt generated by extension of the lithosphere. J. Petrol. 29, 625–679.
- Mei, S., Kohlstedt, D.L., 2000. Influence of water on plastic deformation of olivine aggregates: 1. Diffusion creep regime. J. Geophys. Res. 105, 21457–21469.
- Mei, S., Kohlstedt, D.L., 2000. Influence of water on plastic deformation of olivine aggregates: 2. Dislocation creep regime. J. Geophys. Res. 105, 21471–21481.
- Moresi, L., Solomatov, V., 1998. Mantle convection with a brittle lithosphere: thoughts on the global tectonic styles of the Earth and Venus. Geophys. J. Int. 133, 669–682.
- Nielsen, S.G., Baker, J.A., Krogstad, E.J., 2002. Petrogenesis of an early Archaean (3.4 Ga) norite dyke, Isua, West Greenland: evidence for early Archaean crustal recycling? Precambrian Res. 118, 133–148.
- Nisbet, E.G., 1982. The tectonic setting and petrogenesis of komatiites. In: Arndt, N.T., Nisbet, E.G. (Eds.), Komatiites. George Allen & Unwin, London, pp. 501–520.
- Nisbet, E.G., Cheadle, M.J., Arndt, N.T., Bickle, M.J., 1993. Constraining the potential temperature of the Archaean mantle: a review of the evidence from komatiites. Lithos 30, 291–307.
- Ogawa, M., 1988. Numerical experiments on coupled magmatismmantle convection system: implications for mantle evolution and Archean continental crusts. J. Geophys. Res. 93 (B12), 15119–15134.
- Ogawa, M., 2000. Coupled magmatism–mantle convection system with variable viscosity. Tectonophysics 322, 1–18.
- Parmentier, E.M., Hess, P.C., 1992. Chemical differentiation of a convecting planetary interior: consequences for a one plate planet such as Venus. Geophys. Res. Lett. 19 (20), 2015–2018.
- Patchett, P.J., 1983. Importance of the Lu–Hf isotopic system in studies of planetary chronology and chemical evolution. Geochim. Cosmochim. Acta 47, 81–91.
- Peck, W.J., Valley, J.W., Wilde, S.M., Graham, C.M., 2001. Oxygen isotope ratios and rare earth elements in 3.3 to 4.4 Ga zircons: ion microprobe evidence for high ¹⁸O continental crust and oceans in the Early Archean. Geochim. Cosmochim. Acta 65 (22), 4215–4229.
- Philpotts, A.R., 1990. Principles of Igneous and Metamorphic Petrology. Prentice-Hall, New Jersey.
- Piepenbreier, D., Stöckhert, B., 2001. Plastic flow of omphacite in eclogites at temperatures below 500 °C—implications for interplate coupling in subduction zones. Int. J. Earth Sci. 90, 197–210.
- Presnall, D.C., Weng, Y.-H., Milholland, C.S., Walter, M.J., 1998. Liquidus phase relations in the system MgO-MgSiO₃ at

pressures up to 25 GPa—constraints on crystallization of a molten Hadean mantle. Phys. Earth Planet. Inter. 107, 83–95.

- Ringwood, A.E., Irifune, T., 1988. Nature of the 650-km seismic discontinuity; implications for mantle dynamics and differentiation. Nature 331 (6152), 131–136.
- Rudnick, R.L., Barth, M., Horn, I., McDonough, W.F., 2000. Rutilebearing refractory eclogites: missing link between continents and depleted mantle. Science 287, 278–281.
- Schaber, G.G., Strom, R.G., Moore, H.J., Soderblom, L.A., Kirk, R.L., Chadwick, D.J., Dawson, D.D., Gaddis, L.R., Boyce, J.M., Russel, J., 1992. Geology and distribution of impact craters on Venus: what are they telling us? J. Geophys. Res. 97 (E8), 13257–13301.
- Schott, B., Van den Berg, A.P., Yuen, D.A., 2001. Focussed timedependent martian volcanism from chemical differentiation coupled with variable thermal conductivity. Geophys. Res. Lett. 28 (22), 4271–4274.
- Sleep, N.H., Windley, B.F., 1982. Archean plate tectonics: constrains and inferences. J. Geol. 90, 363–379.
- Solomatov, V.S., Moresi, L.-N., 1996. Stagnant lid convection on Venus. J. Geophys. Res. 101 (E2), 4737–4753.
- Stein, M., Hofmann, A.W., 1994. Mantle plumes and episodic crustal growth. Nature 372, 63–68.
- Steinbach, V., Yuen, D.A., Zhao, W., 1993. Instabilities from phasetransitions and the timescales of mantle thermal evolution. Geophys. Res. Lett. 20 (12), 1119–1122.
- Tackley, P.J., 2000. Mantle convection and plate tectonics: toward an integrated physical and chemical theory. Science 288, 2002–2007.
- Turcotte, D.L., 1995. How does Venus lose heat? J. Geophys. Res. 100 (E8), 16931–16940.
- Van den Berg, A.P., Yuen, D.A., 1998. Modelling planetary dynamics by using the temperature at the core-mantle boundary as a control variable: effects of rheological layering on mantle heat transport. Phys. Earth Planet. Inter. 108, 219–234.
- Van den Berg, A.P., Van Keken, P.E., Yuen, D.A., 1993. The effects of a composite non-Newtonian and Newtonian rheology on mantle convection. Geophys. J. Int. 115, 62–78.
- Van der Hilst, R.D., Kárason, H., 1999. Compositional heterogeneity in the bottom 1000 km of the Earth's mantle: toward a hybrid convection model. Science 283, 1885–1888.
- Van Hunen, J., Van den Berg, A.P., Vlaar, N.J., 2002. On the role of subducting oceanic plateaus in the development of shallow flat subduction. Tectonophysics 352, 317–333.
- Van Keken, P.E., 1997. Evolution of starting mantle plumes: a comparison between numerical and laboratory experiments. Earth Planet. Sci. Lett. 148, 1–11.
- Van Thienen, P., Van den Berg, A.P., De Smet, J.H., Van Hunen, J., Drury, M.R., 2003. Interaction between smallscale mantle diapirs and a continental root. Geochem. Geophys. Geosyst. 4 (doi:10.1029/2002GC000338).
- Vervoort, J.D., Patchett, P.J., Gehrels, G.E., Nutman, A.P., 1996. Constraints on early Earth differentiation from hafnium and neodymium isotopes. Nature 379, 624–627.
- Vlaar, N.J., 1985. Precambrian geodynamical constraints. In: Tobi, A.C., Touret, J.L.R. (Eds.), The Deep Proterozoic Crust in the North Atlantic Provinces. Reidel, Dordecht, pp. 3–20.

24

P. van Thienen et al. / Tectonophysics xx (2004) xxx-xxx

- Vlaar, N.J., 1986. Archaean global dynamics. Geol. Mijnb. 65, 91-101.
- Vlaar, N.J., Van den Berg, A.P., 1991. Continental evolution and archeao-sea-levels. In: Sabadini, R., Lambeck, K., Boschi, E. (Eds.), Glacial Isostasy, Sea-Level and Mantle Rheology. Kluwer Academic Publishing, Dordrecht, Netherlands, pp. 637–662.
- Vlaar, N.J., Van Keken, P.E., Van den Berg, A.P., 1994. Cooling of the Earth in the Archaean: consequences of pressurerelease melting in a hotter mantle. Earth Planet. Sci. Lett. 121, 1–18.
- Walzer, U., Hendel, R., 1997. Time-dependent thermal convection, mantle differentiation and continental crust growth. Geophys. J. Int. 130, 303–325.
- Walzer, U., Hendel, R., 1999. A new convection–fractionation model for the evolution of the principal geochemical reservoirs of the Earth's mantle. Phys. Earth Planet. Inter. 112, 211–256.

- Weinberg, R.F., Podladchikov, Y., 1994. Diapiric ascent of magmas through power law crust and mantle. J. Geophys. Res. 99 (B5), 9543–9559.
- Wilde, S.A., Valley, J.D., Peck, W.H., Graham, C.M., 2001. Evidence from detrital zircons for the existence of continental crust and oceans on the Earth 4.4 Gyr ago. Nature 409, 175–178.
- Yaxley, G.M., Green, D.H., 1998. Reactions between eclogite and peridotite: mantle refertilization by subduction of oceanic crust. Schweiz. Mineral. Petrogr. Mitt. 78, 243–255.
- Zegers, T.E., Van Keken, P.E., 2001. Middle Archean continent formation by crustal delamination. Geology 29, 1083–1086.
- Zhai, M., Zhao, G., Zhang, Q., 2002. Is Dongwanzi complex an Archean Ophiolite? Science 295 (5557), 923.